

# Remote Sensing of Ecosystem Productivity Using MODIS

Fred Huemmrich, UMBC/GSFC



*Goddard*  
SPACE FLIGHT CENTER



# Study Goals

Develop methods using only optical signals to estimate ecosystem carbon exchange

- 1) Examine the relationships between ecosystem production (GEP) and spectral reflectance
  - We have some physical understanding of the nature of these relationships but we do not have a good physical model relating leaf/canopy biochemistry, photosynthetic processes, and spectral reflectance
  - Use data from existing flux towers to empirically examine relationships for different vegetation types over multiple years
- 2) Define an algorithm for a potential MODIS product

# Optical Signals

---

Leaf biochemistry responds to stresses over varying time scales

- Short term stress responses change relative amounts of Xanthophyll cycle pigments in leaves
- There are also longer term changes in the relative amounts of photosynthetic and photoprotective pigments (Chlorophylls and Carotenoids) in leaves

These biochemical changes produce detectable changes in leaf optical properties - we are trying to relate them to carbon fluxes

Using these optical signals as model inputs has an important effect on the interpretation of the model

- We go from trying to predict vegetation response to environmental variables (temperature and humidity)
- To an approach where we are observing the plant's responses to environmental conditions
  - even if we don't know exactly what those environmental forcings are

# Optical Signals

---

The Photochemical Reflectance Index (PRI) is the normalized difference of reflectances at 531 nm (Band 11) and a reference band at 570 nm (which we don't have on MODIS)

- it was developed to detect Xanthophyll pigments
- PRI is also affected by the overall size of the Chlorophyll and Carotenoid pools in leaves
- we are calling the index for this the Chlorophyll-Carotenoid Index (CCI), the normalized difference of bands 11 and 1 (red band)


$$PRI = \frac{(R_{11} - R_{ref})}{(R_{11} + R_{ref})}$$

# Seasonal Change in Boreal Conifer Needles

Time trends for *Pinus contorta* leaves exposed to a boreal climate

Red points - needle photosynthesis

Blue points - chlorophyll:carotenoid ratio



Black line:  
Chlorophyll-Carotenoid Index

$$CCI = \frac{(R_{11} - R_1)}{(R_{11} + R_1)}$$

## Methods

---

### Examined 43 flux tower sites

Chose sites based on visual evaluation that at least 1 km<sup>2</sup> around tower was uniform vegetation type

### Used MAIAC reflectances from Aqua


- Clear observations during growing season
- MODIS bands 1-12 (Land and Ocean bands)
- VZA<45°
- Filtered spikes and removed noisy winter data
- At least two years of data for each site

### Matched MODIS data with daily gross ecosystem productivity (GEP) from LaThuile fluxnet synthesis data


- ~4500 points total

# Study Sites

Site	Site name
AT-Neu	Austria - Neustift/Stubai Valley
BR-Ban	Brazil - Ecotone Bananal Island
BR-Ma2	Brazil - Manaus - ZF2 K34
BR-Sa1	Brazil - Santarem-Km67-Primary Forest
CA-Man	Canada - BOREAS NSA - Old Black Spruce
CA-Oas	Canada - Sask.- SSA Old Aspen
CA-Obs	Canada - Sask.- SSA Old Black Spruce
CA-Ojp	Canada - Sask.- SSA Old Jack Pine
CA-WP1	Canada - Western Peatland- LaBiche
DE-Hai	Germany - Hainich
DE-Meh	Germany - Mehrstedt 1
DE-Tha	Germany - Anchor Station Tharandt - spruce
DE-Wet	Germany - Wetzstein
ES-LMa	Spain - Las Majadas del Tietar
FI-Hyy	Finland - Hyttiala
FR-Pue	France - Puechabon
HU-Bug	Hungary - Bugacpuszta
IT-Col	Italy - Collelongo- Selva Piana
IT-Cpz	Italy - Castelporziano
IT-Lav	Italy - Lavarone (after 3/2002)
IT-MBo	Italy - Monte Bondone
IT-Pia	Italy - Island of Pianosa
IT-Ro1	Italy - Roccarespampani 1
IT-Ro2	Italy - Roccarespampani 2
IT-SRo	Italy - San Rossore
NL-Loo	Netherlands - Loobos
PT-Esp	Portugal - Espirra
PT-Mi1	Portugal - Mitra (Evora)
SE-Nor	Sweden - Norunda
US-Atq	USA - AK - Atqasuk
US-Bar	USA - NH - Bartlett Experimental Forest
US-Ha1	USA - MA - Harvard Forest EMS Tower
US-Ho1	USA - ME - Howland Forest (main tower)
US-Ho2	USA - ME - Howland Forest (west tower)
US-Ivo	USA - AK - Ivotuk
US-LPH	USA - MA - Little Prospect Hill
US-Me2	USA - OR - Metolius-intermediate aged ponderosa pine
US-Me3	USA - OR - Metolius-second young aged pine
US-MMS	USA - IN - Morgan Monroe State Forest
US-NC2	USA - NC - NC_Loblolly Plantation
US-Ne3	USA - NE - Mead - rainfed maize-soybean
US-SRM	USA - AZ - Santa Rita Mesquite
US-Wrc	USA - WA - Wind River Crane Site



# CCI vs. GEP – All Sites



# Analysis Plan

## Evaluate

- What is the best reference band for MODIS PRI?
- How do other vegetation indices perform?
- How does view angle affect relationships?
- How do relationships differ for different vegetation types?
- What is the expected accuracy in retrievals?

Previous studies have looked at these questions but only for a few sites

- Collection 6 processing made it possible to examine lots of sites

# Evaluation of MODIS PRI Reference Bands

	Ref Bands	12	10	1	3	4	
	Site	IGBP class	PRI (11,12)	PRI (11,10)	PRI (11,1)	PRI (11,3)	PRI (11,4)
Deciduous Broadleaf Forest	CA-Oas	DBF		0.03	0.58	0.84	0.49
	DE-Hai	DBF		0.16	0.20	0.82	0.61
	IT-Col	DBF		0.06	0.11	0.86	0.43
	IT-Ro1	DBF		0.05	0.26	0.40	0.39
	IT-Ro2	DBF		0.00	0.17	0.21	0.33
	US-Bar	DBF		0.10	0.56	0.89	0.70
	US-Ha1	DBF		0.03	0.43	0.72	0.55
Evergreen Broadleaf Forest	US-LPH	DBF		0.18	0.63	0.92	0.74
	US-MMS	DBF		0.20	0.55	0.85	0.76
	BR-Ban	EB F		0.01	0.00	0.02	0.00
	BR-Ma2	EB F		0.01	0.00	0.10	0.00
	BR-Sa1	EB F		0.00	0.08	0.06	0.19
	FR-Pue	EB F		0.18	0.00	0.34	0.26
	IT-Cpz	EB F		0.06	0.02	0.18	0.18
Evergreen Needleleaf Forest	PT-Esp	EB F		0.05	0.05	0.00	0.00
	PT-Mi1	EB F		0.10	0.17	0.20	0.23
	CA-Man	EN F		0.07	0.27	0.44	0.05
	CA-Obs	EN F		0.00	0.21	0.65	0.02
	CA-Ojp	EN F		0.11	0.17	0.47	0.00
	DE-Tha	EN F		0.27	0.05	0.59	0.35
	DE-Wet	EN F		0.32	0.17	0.41	0.16
Grassland	FI-Hyy	EN F		0.19	0.03	0.65	0.09
	IT-Lav	EN F		0.45	0.22	0.81	0.80
	IT-SRo	EN F		0.35	0.13	0.01	0.00
	NL-Loo	EN F		0.41	0.30	0.34	0.02
	SE-Nor	EN F		0.00	0.16	0.32	0.04
	US-Ho1	EN F		0.42	0.15	0.78	0.47
	US-Ho2	EN F		0.54	0.01	0.64	0.20
Shrub/Savanna	US-Me2	EN F		0.31	0.19	0.23	0.02
	US-Me3	EN F		0.41	0.31	0.41	0.05
	US-NC2	EN F		0.00	0.58	0.76	0.73
	US-Wrc	EN F		0.21	0.00	0.36	0.07
	AT-Neu	GRA		0.16	0.00	0.49	0.31
	DE-Meh	GRA		0.05	0.35	0.65	0.57
	HU-Bug	GRA		0.16	0.15	0.02	0.01
Wetlands	IT-MBo	GRA		0.00	0.16	0.82	0.54
	IT-Pia	OSH		0.25	0.29	0.44	0.39
	ES-LMa	SAV		0.01	0.19	0.32	0.25
	US-SRM	WSA		0.01	0.15	0.20	0.05
	US-Ne3	CRO		0.10	0.43	0.47	0.49
	US-Atq	WET		0.08	0.10	0.10	0.12
	US-Ivo	WET		0.12	0.01	0.00	0.00
	CA-WP1	WET (MF)		0.05	0.68	0.81	0.54

Table shows R<sup>2</sup> with Daily GEP  
Color coded as:  
R<sup>2</sup> ≥ 0.5 - Yellow  
R<sup>2</sup> ≥ 0.7 - Red

$$PRI(11, ref) = \frac{(R_{11} - R_{ref})}{(R_{11} + R_{ref})}$$

PRI(11,1) = Chlorophyll-Carotenoid Index (CCI)

# Evaluation of Other MODIS Vegetation Indices

Site	IGBP class	NDVI	EVI	Chl	NDWI	NDII	PRI (11,1)
CA-Oas	DBF	0.78	0.80	0.75	0.42	0.57	0.84
DE-Hai	DBF	0.83	0.88	0.73	0.78	0.11	0.82
IT-Col	DBF	0.79	0.84	0.54	0.63	0.03	0.86
IT-Ro1	DBF	0.58	0.72	0.60	0.61	0.01	0.40
IT-Ro2	DBF	0.46	0.60	0.60	0.45	0.01	0.21
US-Bar	DBF	0.84	0.89	0.45	0.82	0.53	0.89
US-Ha1	DBF	0.58	0.68	0.46	0.69	0.56	0.72
US-LPH	DBF	0.83	0.91	0.67	0.85	0.79	0.92
US-MMS	DBF	0.78	0.87	0.74	0.85	0.77	0.85
BR-Ban	EB F	0.00	0.05	0.00	0.00	0.00	0.02
BR-Ma2	EB F	0.01	0.01	0.00	0.00	0.01	0.10
BR-Sa1	EB F	0.01	0.20	0.03	0.15	0.04	0.06
FR-Pue	EB F	0.00	0.32	0.04	0.01	0.00	0.34
IT-Cpz	EB F	0.02	0.22	0.01	0.20	0.00	0.18
PT-Esp	EB F	0.03	0.00	0.04	0.06	0.01	0.00
PT-Mi1	EB F	0.24	0.33	0.27	0.31	0.03	0.20
CA-Man	EN F	0.11	0.54	0.01	0.00	0.00	0.44
CA-Obs	EN F	0.26	0.46	0.00	0.12	0.21	0.65
CA-Ojp	EN F	0.08	0.06	0.23	0.14	0.00	0.47
DE-Tha	EN F	0.48	0.58	0.15	0.19	0.01	0.59
DE-Wet	EN F	0.07	0.36	0.00	0.20	0.00	0.41
FI-Hyy	EN F	0.46	0.47	0.24	0.09	0.00	0.65
IT-Lav	EN F	0.65	0.62	0.17	0.08	0.02	0.81
IT-SRo	EN F	0.08	0.03	0.05	0.20	0.00	0.01
NL-Loo	EN F	0.03	0.43	0.00	0.09	0.00	0.34
SE-Nor	EN F	0.45	0.17	0.51	0.24	0.13	0.32
US-Ho1	EN F	0.46	0.63	0.06	0.02	0.04	0.78
US-Ho2	EN F	0.18	0.44	0.02	0.00	0.03	0.64
US-Me2	EN F	0.38	0.01	0.38	0.34	0.09	0.23
US-Me3	EN F	0.28	0.01	0.37	0.31	0.26	0.41
US-NC2	EN F	0.64	0.76	0.48	0.72	0.65	0.76
US-Wrc	EN F	0.06	0.15	0.05	0.10	0.06	0.36
AT-Neu	GRA	0.41	0.41	0.24	0.52	0.00	0.49
DE-Meh	GRA	0.71	0.74	0.69	0.73	0.06	0.65
HU-Bug	GRA	0.09	0.03	0.14	0.05	0.04	0.02
IT-MBo	GRA	0.76	0.79	0.52	0.72	0.05	0.82
IT-Pia	OSH	0.45	0.44	0.38	0.44	0.02	0.44
ES-LMa	SAV	0.32	0.46	0.30	0.32	0.03	0.32
US-SRM	WSA	0.61	0.72	0.41	0.45	0.19	0.20
US-Ne3	CRO	0.64	0.66	0.62	0.63	0.56	0.47
US-Atq	WET	0.24	0.10	0.17	0.04	0.07	0.10
US-Ivo	WET	0.03	0.03	0.06	0.10	0.03	0.00
CA-WP1	WET (MF)	0.47	0.78	0.01	0.28	0.18	0.81

Table shows R<sup>2</sup> with Daily GEP

Color coded as:  
R<sup>2</sup> ≥ 0.5 - Yellow  
R<sup>2</sup> ≥ 0.7 - Red

$$NDVI = \frac{(R_2 - R_1)}{(R_2 + R_1)}$$

$$EVI = \frac{2.5 * (R_2 - R_1)}{(R_2 + 6 * R_1 + 7.5 * R_3 + 1)}$$

$$Chl = \left[ \frac{1}{R_1} - \frac{1}{R_2} \right] * R_2$$

$$NDWI = \frac{(R_2 - R_5)}{(R_2 + R_5)}$$

$$NDII = \frac{(R_2 - R_6)}{(R_2 + R_6)}$$

$$PRI(11,1) = \frac{(R_{11} - R_1)}{(R_{11} + R_1)}$$





# View Angle Effect on CCI

		F	B	N	
		All Pts. R2 CCI	Forward R2 CCI	Back R2 CCI	
	Site ID	IGBP class			
Deciduous Broadleaf Forest	CA-Oas	DBF	0.84	0.87	0.85
	DE-Hai	DBF	0.82	0.78	0.80
	IT-Col	DBF	0.86	0.82	0.89
	IT-Ro1	DBF	0.40	0.29	0.38
	IT-Ro2	DBF	0.21	0.13	0.21
	US-Bar	DBF	0.89	0.93	0.90
	US-Ha1	DBF	0.72	0.71	0.75
	US-LPH	DBF	0.92	0.91	0.94
	US-MMS	DBF	0.85	0.82	0.84
	BR-Ban	EBF	0.02	0.03	0.01
Evergreen Broadleaf Forest	BR-Ma2	EBF	0.10	0.26	0.03
	BR-Sa1	EBF	0.06	0.03	0.00
	FR-Pue	EBF	0.34	0.51	0.26
	IT-Cpz	EBF	0.18	0.21	0.24
	PT-Esp	EBF	0.00	0.01	0.09
	PT-Mi1	EBF	0.20	0.26	0.14
	CA-Man	ENF	0.44	0.47	0.53
	CA-Obs	ENF	0.65	0.65	0.64
	CA-Ojp	ENF	0.47	0.53	0.48
	DE-Tha	ENF	0.59	0.62	0.69
Evergreen Needleleaf Forest	DE-Wet	ENF	0.41	0.55	0.52
	FI-Hyy	ENF	0.65	0.80	0.73
	IT-Lav	ENF	0.81	0.88	0.89
	IT-SRo	ENF	0.01	0.10	0.02
	NL-Loo	ENF	0.34	0.57	0.12
	SE-Nor	ENF	0.32	0.54	0.33
	US-Ho1	ENF	0.78	0.83	0.82
	US-Ho2	ENF	0.64	0.71	0.75
	US-Me2	ENF	0.23	0.34	0.27
	US-Me3	ENF	0.41	0.56	0.46
Grassland	US-NC2	ENF	0.76	0.77	0.76
	US-Wrc	ENF	0.36	0.48	0.54
	AT-Neu	GRA	0.49	0.46	0.54
	DE-Meh	GRA	0.65	0.67	0.78
Shrub/Savanna	HU-Bug	GRA	0.02	0.06	0.07
	IT-MBo	GRA	0.82	0.89	0.81
	IT-Pia	OSH	0.44	0.45	0.57
Wetlands	ES-LMa	SAV	0.32	0.40	0.25
	US-SRM	WSA	0.20	0.29	0.26
	US-Ne3	CRO	0.47	0.48	0.47
	CA-WP1	WET	0.81	0.82	0.89
Wetlands	US-Atq	WET	0.10	0.33	0.06
	US-Ivo	WET	0.00	0.01	0.00

Table shows R<sup>2</sup> with Daily GEP  
 - Nadir - All VAZ when  
 VZA<15°


Highlighted cell has highest R<sup>2</sup>  
 - Red means difference  
 between max and min R<sup>2</sup> >0.15

# CCI and EVI vs. Daily GEP for all sites combined




# CCI vs. Daily GEP – Grouped by IGBP Class


DBF




EBF




ENF




GRA



OSH, SAV, WSA




WET




# EVI vs. Daily GPP – Grouped by IGBP Class


DBF




EBF




ENF




GRA



OSH, SAV, WSA




WET




# Multiple Regression of MODIS Bands on Daily GEP


DBF




EBF




ENF




GRA



OSH, SAV, WSA





WET




Used all bands except band 6

# CCI and Multiple Regression vs. Daily GEP for all sites combined



# Multiple Regression Coefficients



# Conclusions

---

- Although not designed for this purpose, MODIS reflectances combining land and ocean bands can be used to derive GEP
  - Need more training data to develop robust algorithm
  - Often hard to find tower sites with uniform vegetation covering a large enough area, particularly for some vegetation types (e.g. crops)
- CCI and EVI appear to be the most promising vegetation indices
  - Multiple regressions using all bands works even better
- Effects of view angle can be significant, but presently cannot predict effect
  - variations may be more related to spatial distribution of vegetation
- RMSE of retrievals of daily GEP on the order of  $\sim 2 \text{ gC m}^{-2} \text{ d}^{-1}$
- There are different relationships at different sites
  - Stratifying by vegetation type can help some

# Thanks

---

Alexei Lyapustin and Yujie Wang  
Dave Landis  
John Gamon

## Fluxnet and all of the flux providers

This work used eddy covariance data acquired by the FLUXNET community and in particular by the following networks: AmeriFlux (U.S. Department of Energy, Biological and Environmental Research, Terrestrial Carbon Program (DE-FG02-04ER63917 and DE-FG02-04ER63911)), CarboEuropeIP, CarboItaly, CarboMont, Fluxnet-Canada (supported by CFCAS, NSERC, BIOCAP, Environment Canada, and NRCan), GreenGrass, LBA, NECC, USCCC. We acknowledge the financial support to the eddy covariance data harmonization provided by CarboEuropeIP, FAO-GTOS-TCO, iLEAPS, Max Planck Institute for Biogeochemistry, National Science Foundation, University of Tuscia, Université Laval, Environment Canada and US Department of Energy and the database development and technical support from Berkeley Water Center, Lawrence Berkeley National Laboratory, Microsoft Research eScience, Oak Ridge National Laboratory, University of California – Berkeley and the University of Virginia.